arXiv:1704.03551v1 [math.AG] 11 Apr 2017

ON THE MULTICANONICAL SYSTEMS OF QUASI-ELLIPTIC SURFACES IN CHARACTERISTIC 3

TOSHIYUKI KATSURA

ABSTRACT. We consider the multicanonical systems $|mK_S|$ of quasi-elliptic surfaces with Kodaira dimension 1 in characteristic 3. We show that for any $m \geq 5$ $|mK_S|$ gives the structure of quasi-elliptic fiber space, and the number 5 is best possible to give the structure for any such surfaces.

Dedicated to Professor Piotr Pragacz on the occasion of his sixtieth birthday

1. INTRODUCTION

Let k be an algebraically closed field in characteristic $p > 0$, and let $\varphi : S \longrightarrow B$ be an elliptic surface with Kodaira dimension $\kappa(S) = 1$ over k. Let K_S be a canonical divisor of S. We consider the multicanonical system $|mK_S|$. In Katsura-Ueno [4] and Katsura [3], we considered the following question:

Question 1.1. Is there a positive integer M such that if $m \geq M$, then the multicanonical system $|mK_S|$ gives a structure of elliptic surface for any elliptic surface S over k with $\kappa(S) = 1$?

For the complex analytic case, Iitaka showed that $M = 86$ and 86 is best possible (cf. Iitaka [2]). Namely, if m is smaller than 86, then there exists an elliptic surface $\varphi : S \longrightarrow B$ with $\kappa(S) = 1$ such that $|mK_S|$ does not give the structure of elliptic surface. In the case of algebraic elliptic surfaces, if the characteristic $p = 0$ or $p \geq 3$, we showed that $M = 14$ and 14 is best possible (Katsura-Ueno [4] and Katsura [3]). If $p = 2$, then we showed that $M = 12$ and 12 is best possible (cf. Katsura [3]).

In this paper, we treat quasi-elliptic surfaces $\varphi : S \longrightarrow B$ and consider the following similar question.

Question 1.2. Is there a positive integer M such that if $m \geq M$, then the multicanonical system $|mK_S|$ gives the structure of quasi-elliptic surface for any quasi-elliptic surface S over k with $\kappa(S) = 1$?

Note that quasi-elliptic surfaces exist only in characterisitics $p = 2$ and 3 (cf. Bombieri-Mumford [1]). We show that if $p = 3$, then we have $M = 5$ and the number 5 is best

Research of the author is partially supported by Grant-in-Aid for Scientific Research (B) No. 15H03614.

possible. In characteristic $p = 2$, we have still some difficulties to determine the best possible number.

In Section 2, we summarize basic facts on quasi-elliptic surfaces. In Section 3, we examine the multicanonical system of quasi-elliptic surfaces in characteristic 3 and show our main theorem.

2. Some lemmas for quasi-elliptic surfaces

Let k be an algebraically closed field in characteristic $p > 0$ and let $\varphi : S \longrightarrow B$ be a quasi-elliptic surface defined over k. Throughout this paper, we assume that any exceptional curve of the first kind is not contained in fibers. Such a surface exists if and only if p is equal to 2 or 3, and the multiplicities of multiple fibers are all equal to p (cf. Bombieri-Mumford [1]). We denote by g the genus of the curve B. In this section, we recall some facts on quasi-elliptic surfaces. We denote by pF_i $(i = 1, \ldots, \lambda)$ the multiple fibers. Let $\mathcal{T}$ be the torsion part of $R^1\varphi_*\mathcal{O}_S$. Then, there exists a Cartier divisor $\mathbf{f}$ on B such that $R^1\varphi_*\mathcal{O}_S/\mathcal{T} \cong \mathcal{O}_B(\mathbf{f})$. The canonical divisor formula of S is given by

$$K_S \sim \varphi^*(K_B - \mathbf{f}) + \sum_{i=1}^{\lambda} a_i F_i,$$

where $-\deg\mathbf{f} = \chi(\mathcal{O}_S) + t$ with $t =$ the length of the torsion part of $R^1\varphi_*\mathcal{O}_S$ and $0 \leq a_i \leq p-1$. Here, $\sim$ means linear equivalence. If pF_i is a tame multiple fiber, then we have $a_i = p - 1$. For details, see Bombieri-Mumford [1].

Lemma 2.1. *The Albanese variety* $\mathrm{Alb}(S)$ *of* S *is isomorphic to the Jacobian variety* $\mathrm{J}(B)$ *of* B.

Proof. Let $\psi : S \longrightarrow \mathrm{Alb}(S)$ be the Albanese mapping. If $\mathrm{Alb}(S)$ is a point, then by the universality of Albanese variety we see that the Jacobian variety $\mathrm{J}(B)$ of B is also a point. Now, assume $\mathrm{Alb}(S)$ is not a point. Since the general fiber of φ is a rational curve with one cusp, the fibers are contracted by ψ. Therefore, $\psi(S)$ is a curve. We have, by the universality of Albanese variety, a commutative diagram:

$$\begin{array}{rccc} \varphi : & S & \longrightarrow & B \\ & \psi\downarrow & & \downarrow \\ & \mathrm{Alb}(S) & \longrightarrow & \mathrm{J}(B) \\ & \cup & & \cup \\ & \psi(S) & & B. \end{array}$$

By this diagram, we have a morphism $\psi(S) \longrightarrow B$. Therefore, by the Stein factorization theorem, we see that $\psi(S)$ is isomorphic to B. Therefore, by the universality of Jacobian variety, we conclude $\mathrm{Alb}(S) \cong \mathrm{J}(B)$ (see also Katsura-Ueno [4], Lemma 3.4).

We find the following lemma and corollary in Lang [5] and Raynaud [6]. We give here an easy proof for the lemma.

Lemma 2.2. *Let $\varphi : S \longrightarrow B$ be a quasi-elliptic surface over a non-singular complete curve B with genus g. Then, we have the inequality $\chi(\mathcal{O}_S) \geq (1-g)/3$.*

Proof. By Noether's formula and the self-intersection number $c_1(S)^2 = 0$ of the first Chern class of S, we have

$$12\chi(\mathcal{O}_S) = c_1(S)^2 + c_2(S) = 2 - 4q(S) + b_2(S).$$

By Lemma 2.1, we have $q(S) = g$. Denoting by $\rho(S)$ the Picard number of S, we have also $b_2(S) \geq \rho(S) \geq 2$. Hence, we have $\chi(\mathcal{O}_S) \geq (1-g)/3$.

Corollary 2.3. (i) *If $g = 1$, then $\chi(\mathcal{O}_S) \geq 0$.*
(ii) *If $g = 0$, then $\chi(\mathcal{O}_S) \geq 1$.*

3. Multicanonical systems

In this section, let k be an algebraically closed field of characteristic 3. Let $\varphi : S \longrightarrow B$ be a quasi-elliptic surface defined over k.

Example 3.1. In characteristic 3, we consider the quasi-elliptic surface $\varphi : S \longrightarrow \mathbf{P}^1$ which is given by a non-singular complete model of the surface defined by

$$t^2(t-1)^2 z + t^2(t-1)^2 + x^3 + tz^3 = 0$$

Here, t is a parameter of the base curve $\mathbf{P}^1$. By Lang [5] p.485, this surface has two tame multiple fibers at $t = 0, 1$, and we have $\chi(\mathcal{O}_S) = 1$. We denote the two tame multiple fibers by $3F_0$ and $3F_1$. The canonical divisor K_S is given by

$$\begin{aligned} K_S &\sim \varphi^*(K_{\mathbf{P}^1} - \mathbf{f}) + 2F_0 + 2F_1 \\ &\sim -F + 2F_0 + 2F_1. \end{aligned}$$

Here, $\mathbf{f}$ is a Cartier divisor on $\mathbf{P}^1$ with $-\mathrm{deg}\mathbf{f} = \chi(\mathcal{O}_S) = 1$ such that $\mathcal{O}_{\mathbf{P}^1}(\mathbf{f}) \cong R^1\varphi_*\mathcal{O}_S$, and F is a general fiber of $\varphi : S \longrightarrow \mathbf{P}^1$. Since we have $F \sim 3F_0 \sim 3F_1$, we see $4K_S \sim 2F_0 + 2F_1$. Therefore, we have $\dim \mathrm{H}^0(S, \mathcal{O}_S(4K_S)) = 1$, and $|4K_S|$ does not give the structure of quasi-elliptic surface. If $m \geq 5$, then we have $\dim \mathrm{H}^0(S, \mathcal{O}_S(mK_S)) \geq 2$, and $|mK_S|$ gives the structure of quasi-elliptic surface.

We have the following theorem.

Theorem 3.2. *Assume that the characteristic $p = 3$. Then, for any quasi-elliptic surface $f : S \to B$ with $\kappa(S) = 1$ over k and for any $m \geq 5$, the multicanonical system $|mK_S|$ gives the unique structure of quasi-elliptic surface, and the number 5 is best possible.*

Proof. The method of the proof is similar to the one in Iitaka [2], Katsura-Ueno [4] and Katsura [3]. Since the Kodaira dimension is equal to 1, the structure of quasi-elliptic

surface is unique. The Kodaira dimension of S is equal to 1 if and only if

$$(*) \quad 2g-2+\chi(\mathcal{O}_S)+t+\sum_{i=1}^{\lambda}(a_i/3)>0.$$

Therefore, we need to find the least integer m such that

$$(**) \quad m(2g-2+\chi(\mathcal{O}_S)+t)+\sum_{i=1}^{\lambda}[ma_i/3]\geq 2g+1$$

holds under the condition $(*)$. Here, $[r]$ means the integral part of a real number r. We have the following 6 cases:

Case (I) $g\geq 2$
Case (II-1) $g=1, \chi(\mathcal{O}_S)+t\geq 1$
Case (II-2) $g=1, \chi(\mathcal{O}_S)=0, t=0$
Case (III-1) $g=0, \chi(\mathcal{O}_S)+t\geq 3$
Case (III-2) $g=0, \chi(\mathcal{O}_S)+t=2$
Case (III-3) $g=0, \chi(\mathcal{O}_S)=1, t=0$

We check $(**)$ under the condition $(*)$ for each case. In Case (I), by Lemma 2.2, we have $2g-2+\chi(\mathcal{O}_S)\geq 5(g-1)/3$. Hence, if $m\geq 3$, $(**)$ holds. In Case (II-1), if $m\geq 3$, $(**)$ holds. In Case (II-2), all multiple fibers are tame in this case, and we have at least one multiple fiber by $(*)$. Since $a_i=2$, $(**)$ holds for $m\geq 5$. In Case (III-1), $(**)$ holds for $m\geq 1$. In Case (III-2), since $\chi(\mathcal{O}_S)\geq 1$ by Corollary 2.3, we have $t\leq 1$. Therefore, the number of wild fibers is less than or equal to 1. If there exists at least one tame multiple fiber, then $(**)$ holds for $m\geq 2$. If there exist no tame fibers and only one wild fiber, then by $(*)$ we have $a_1\geq 1$. Therefore, $(**)$ holds for $m\geq 3$. In Case (III-3), all multiple fibers are tame, and we have $\lambda\geq 2$ by $(*)$. Therefore, $(**)$ holds for $m\geq 5$. The result on the best possible number in characteristic 3 follows from Example 3.1.

In characteristic 2, we can also consider a similar question to the one in characteristic 3. We have still difficulties to decide the best possible number. For example we need to solve the following question.

Question 3.3. Does there exist a quasi-elliptic surface over an elliptic curve with only one tame multiple fiber and with $\chi(\mathcal{O}_S)=0$ in characteristic 2?

If there don't exist such quasi-elliptic surfaces, then we can show that in characteristic 2, $(**)$ holds for $m\geq 4$ and that the best possible number is equal to 4. Namely, we have $M=4$ in characteristic 2.

REFERENCES

[1] E. Bombieri and D. Mumford, Enriques' classification of surfaces in char. p, II, In *Complex Analysis and Algebraic Geometry* (eds. by W.L. Baily, Jr., and T. Shioda), Iwanami Shoten, Tokyo, and Princeton Univ. Press, Princeton, NJ, 1977, 22-42.

[2] S. Iitaka, Deformations of compact complex surfaces, II, *J. Math. Soc. Japan*, **22** (1970), 247-261.
[3] T. Katsura, Multicanonical systems of elliptic surfaces in small characteristics, *Compositio Math.*, **97** (1995), 119-134.
[4] T. Katsura and K. Ueno, On elliptic surfaces in characteristic p, *Math. Ann.*, **272** (1985), 291-330.
[5] W. Lang, Quasi-elliptic surfaces in characteristic three, *Ann. Scient. Ec. Norm. Sup.*, **12** (1979), 473-500.
[6] M. Raynaud, Surfaces elliptiques et quasi-elliptiques, manuscript.

FACULTY OF SCIENCE AND ENGINEERING, HOSEI UNIVERSITY, KOGANEI-SHI, TOKYO 184-8584, JAPAN

E-mail address: toshiyuki.katsura.tk@hosei.ac.jp